

BLOAT GUARD- poloxalene powder
Phibro Animal Health

Bloat Guard Top Dressing for Feed® (poloxalene)

Medicated Top Dressing for Feed

For prevention of legume (alfalfa, clover) and wheat pasture bloat in cattle

Active Ingredient

Poloxalene 53%
(Each 2/3 oz by weight of top dressing contains 10 g of poloxalene)

Inert Ingredient:

Verxite® #4..... 47%

CAUTION:

For use in manufacturing medicated animal feeds only.

CAUTION:

Certain components of animal feeds, including medicated premixes, possess properties that may be a potential health hazard or a source of personal discomfort to certain individuals who are exposed to them. Human exposure should, therefore, be minimized by observing the general industry standards for occupational health and safety.

Precautions such as the following should be considered: dust masks or respirators and protective clothing should be worn; dust arresting equipment and adequate ventilation should be utilized; personal hygiene should be observed; wash before eating or leaving a work site; be alert for signs of allergic reactions—seek prompt medical treatment if such reactions are suspected.

STORE AT ROOM TEMPERATURE

Directions for Use as Top Dress on Feed

Use the enclosed measure which is equal to 1/4 of a standard measuring cup and holds approximately 2/3 oz of Bloat Guard by weight. Each 2/3 oz by weight contains 10 g of poloxalene (the active drug ingredient).

Daily Dosage

Animal Weight (lb) Number of Measures

Less than 500	1/2–1
500–1000	1–2
1000–1500	1 1/2–3
Over 1500	2–4

Bloat Guard is to be consumed daily. Bloat Guard should be fed as a top dressing on individual rations of ground feed, starting 2 or 3 days before animals are exposed to bloat-producing conditions. The higher dosage levels are recommended when bloat-producing conditions are severe.

Repeat the feeding of Bloat Guard when animals are exposed to bloat-producing conditions more than 12 hours from the last feeding of Bloat Guard, but do not exceed the higher dosage levels in any 24-hour period.

If your animals do not accept Bloat Guard readily, stir the recommended amount into their feed. After animals become accustomed to the change in diet, use Bloat Guard as a top dressing.

Caution: It is essential that each animal consumes the total recommended dosage of Bloat Guard daily for adequate protection.

Warning:

The normal life of this product is at least 24 months. However, when the product is subjected to extreme temperatures (38°C/100°F) for long periods of time (6 months), spontaneous combustion may occur. The product is not combustible unless it develops a strong, irritating odor; if this occurs, flush with water and discard immediately.

NOT FOR HUMAN USE

Follow Directions on Back Panel

Bloat Guard® is a Phibro Animal Health registered trademark for Poloxalene

Phibro Animal Health, Teaneck, NJ 07666.

Verxite® is a registered trademark of

W.R. Grace & Company.

6080000

101-9030-07



Bloat Guard®

(poloxalene)

Indications for Use:
For prevention of legume (alfalfa, clover) and wheat pasture bloat in cattle



Directions for Use

The dose of Bloat Guard is proportional to body weight and also depends upon the severity of the bloat-producing conditions. The dose of 1 g of poloxalene per 100 lb of body weight is recommended for cattle under moderate bloat-producing conditions. For cattle under severe bloat producing conditions, the dose should be doubled (2 g of poloxalene per 100 lb of body weight).

Use the enclosed measure which is equal to 1/4 of a standard measuring cup and holds approximately 2/3 oz of Bloat Guard by weight. Each 2/3 oz by weight contains 10 g of poloxalene (the active drug ingredient).

Daily Dose

Animal weight (lb)	Number of measures	Dose (grams)
500	1/2 to 1	5 to 10
1000	1 to 2	10 to 20
1500	1 1/2 to 3	15 to 30
2000	2 to 4	20 to 40

Bloat Guard is to be consumed daily. Bloat Guard should be sprinkled on individual rations of ground feed, starting 2 or 3 days before animals are exposed to bloat-producing conditions.

Repeat the feeding of Bloat Guard when animals are exposed to bloat-producing conditions more than 12 hours from the last feeding of Bloat Guard, but do not exceed the higher dose level in any 24-hour period.

If your animals do not accept Bloat Guard readily, stir the recommended amount into their feed. After animals become accustomed to the change in diet, sprinkle Bloat Guard on top of the feed.

Caution: It is essential that each animal consumes the total recommended dose of Bloat Guard daily for adequate protection.

Warning: The normal life of this product is at least 24 months. However, when the product is subjected to extreme temperatures (38°C/100°F) for long periods of time (6 months), spontaneous combustion may occur. The product is not combustible unless it develops a strong, irritating odor; if this occurs, flush with water and discard immediately.

Vantix® is a registered trademark of W.R. Grace & Company.
Store At Or Below 25°C (77°F). Excursions Permitted Up To 40°C (104°F)
Not For Human Use

6080000
101-9030-08

Distributed by
Phibro Animal Health Corporation
Tussock, NJ 07063

Bloat Guard®
(poloxalene)



Bloat Guard®

(poloxalene)

Indications for Use:
For prevention of legume (alfalfa, clover) and wheat pasture bloat in cattle

Active Ingredient:

Poloxalene 53%
(Each 2/3 oz by weight contains 10 g of poloxalene)

Inert Ingredient:

Ventix® #4 47%

CAUTION: Certain components of animal feeds, including medicated pre-mixes, possess properties that may be a potential health hazard or a source of personal discomfort to certain individuals who are exposed to them. Human exposure should, therefore, be minimized by observing the general industry standards for occupational health and safety.

Precautions such as the following should be considered: dust masks or respirators and protective clothing should be worn; dust arresting equipment and adequate ventilation should be utilized; personal hygiene should be observed; wash before eating or leaving a work site; be alert for signs of allergic reactions—seek prompt medical treatment if such reactions are suspected.

Follow Directions on Back Panel

Bloat Guard® is a Phibro Animal Health Corporation registered trademark for Poloxalene.
Phibro Animal Health Corporation, Tussock, NJ 07065.



NADA #32-704, Approved by FDA
6080000
101-9030-08

Net Weight 20 lb (9.08 kg)



BLOAT GUARD

poloxalene powder

Product Information

Product Type	OTC TYPE A MEDICATED ARTICLE ANIMAL DRUG	Item Code (Source)	NDC:66 104-6080
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POLOXALENE (UNII: V8B3K56SW0) (POLOXALENE - UNII:V8B3K56SW0)	POLOXALENE	240.4 g in 0.45 kg

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66104-6080-0	9.08 kg in 1 BAG		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA032704	03/10/1966	

Labeler - Phibro Animal Health (006989008)

Registrant - Phibro Animal Health (006989008)

Revised: 1/2019

Phibro Animal Health